



ehponline.org

ENVIRONMENTAL HEALTH PERSPECTIVES

Variability of Organophosphorous Pesticide Metabolite Levels in Spot and 24-hr Urine Samples Collected from Young Children during 1 Week

**Asa Bradman, Katherine Kogut, Ellen A. Eisen,
Nicholas P. Jewell, Lesliam Quirós-Alcalá, Rosemary Castorina,
Jonathan Chevrier, Nina T. Holland, Dana Boyd Barr,
Geri Kavanagh-Baird, Brenda Eskenazi**

<http://dx.doi.org/10.1289/ehp.1104808>

Online 9 October 2012



NIEHS

National Institute of
Environmental Health Sciences

National Institutes of Health
U.S. Department of Health and Human Services

**Variability of Organophosphorous Pesticide Metabolite Levels in Spot and 24-hr Urine
Samples Collected from Young Children during 1 Week**

Asa Bradman*¹, Katherine Kogut*¹, Ellen A. Eisen^{1,2}, Nicholas P. Jewell^{1,3}, Lesliam Quirós-Alcalá¹, Rosemary Castorina¹, Jonathan Chevrier¹, Nina T. Holland¹, Dana Boyd Barr⁴, Geri Kavanagh-Baird¹, Brenda Eskenazi¹

*A. Bradman and K. Kogut share lead authorship.

¹Center for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, California, USA

²Division of Environmental Health Sciences, School of Public Health, University of California, Berkeley, California, USA

³Division of Biostatistics, School of Public Health, University of California, Berkeley, California, USA

⁴Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, USA

Institutions and addresses where work was performed:

University of California, Berkeley, School of Public Health
Center for Environmental Research and Children's Health (CERCH)
1995 University Ave, Ste 265
Berkeley, CA 94704

CHAMACOS Project
Natividad Medical Center
1441 Constitution Blvd, Bldg 830
Salinas, CA 93906

Centers for Disease Control and Prevention
1600 Clifton Rd
Atlanta, GA 30333

Corresponding author:

Asa Bradman, PhD
Center for Environmental Research and Children's Health (CERCH)
School of Public Health
University of California, Berkeley
1995 University Avenue, Suite 265
Berkeley, CA 94704
Phone: (510) 643-3023 Fax: (510) 642-9083
abradman@berkeley.edu

Short Running title: Variability of pesticide metabolites in children

Keywords: Biomarkers, children, exposure, metabolites, organophosphorous, pesticides, urine, variability

Acknowledgments: This research was supported by grant numbers RD 83171001 and RD 876709 from EPA, and PO1 ES009605 from NIEHS. Its contents are solely the responsibility of the authors and do not necessarily represent the views of the EPA or the NIEHS. We thank the anonymous reviewers for insightful comments and suggestions to improve the paper. We thank Suril Mehta and Todd Whitehead for statistical assistance and also thank our staff, community partners, and study participants.

Competing Financial Interests: B. Eskenazi recently consulted on a pesticide exposure case. Other authors declare no competing financial interests.

Abbreviations:

AIC: Akaike Information Criterion

DAP: Dialkyl phosphate

DE: Diethyl phosphate

DEDTP: Diethyldithiophosphate

DEP: Diethylphosphate

DETP: Diethylthiophosphate

DF: Detection frequency

DM: Dimethyl phosphate

DMDTP: Dimethyldithiophosphate

DMP: Dimethylphosphate

DMTP: Dimethylthiophosphate

FMV: First morning void

GM: Geometric mean

ICC: Intraclass correlation coefficient

LOD: Limit of detection

OP: Organophosphorous

SD: Standard Deviation

UER: Urinary Excretion Rate

ABSTRACT

Background: Dialkyl phosphate (DAP) metabolites in spot urine samples are frequently used to characterize children's exposures to organophosphorous (OP) pesticides. However, variable exposure and short biological half-lives of OP pesticides could result in highly variable measurements, leading to exposure misclassification.

Objective: To examine within- and between-child variability in DAP metabolites in urine samples collected during one week.

Methods: We collected spot urine samples over seven consecutive days from 25 children (ages 3-6 years). On two of the days, we collected 24-hour voids. We assessed the reproducibility of urinary DAP metabolite concentrations and evaluated the sensitivity and specificity of spot urine samples as predictors of high (top 20%) or elevated (top 40%) weekly average DAP metabolite concentrations.

Results: Within-child variance exceeded between-child variance by a factor of two to eight, dependent on metabolite grouping. Though total DAP concentrations in single spot urine samples were moderately to strongly associated with concentrations in same-day 24-hour samples ($r \approx 0.6-0.8$, $p < 0.01$), concentrations in spot samples collected more than one day apart and in 24-hour samples collected three days apart were weakly correlated ($r \approx -0.21-0.38$). Single spot samples predicted high (top 20%) and elevated (top 40%) full-week average total DAP excretion with only moderate sensitivity (≈ 0.52 and ≈ 0.67 , respectively) but relatively high specificity (≈ 0.88 and ≈ 0.78 , respectively).

Conclusions: The high variability we observed in children's DAP metabolite concentrations suggests that single-day urine samples provide only a brief snapshot of exposure. Sensitivity analyses suggest that classification of cumulative OP exposure based on spot samples is prone to type II classification errors.

Introduction

Spot urine samples have been used to assess organophosphorous (OP) pesticide exposures in epidemiological and biomonitoring studies, with samples analyzed for either non-specific dialkyl phosphate (DAP) metabolites (Centers for Disease Control and Prevention 2003) or pesticide-specific metabolites (Fenske et al. 2002). Because collection of spot urine samples is relatively simple and noninvasive, this sample type is commonly used in studies involving children (Fenske et al. 2005). Several studies suggest that spot samples provide meaningful measures of exposure, reporting associations, for example, between residence near farmlands and OP metabolite levels in children's urine (Fenske et al. 2000). However, concerns have been raised as to whether metabolites measured in a single spot sample accurately reflect exposure for the full day or beyond.

Several factors may result in exposure misclassification when using spot samples: (1) Because the concentration of a chemical, urine volume, and rate of urinary excretion vary with fluid and salt intake, time of day, and other factors (Barr et al. 2005; Boeniger et al. 1993; Cornelis et al. 1996), exposure measurements may be more variable than other sample types, such as first morning voids (FMVs) (Kissel et al. 2005), which are more concentrated and reflect a longer period of accumulation, or 24-hour urine collections, considered by some as the “gold standard” (Aprea et al. 2002; Boeniger et al. 1993; Fenske and Elkner 1990; Martin et al. 1996); (2) due to within-child variability in creatinine excretion (Barr et al. 2005; Boeniger et al. 1993), creatinine adjustment may introduce new variability that does not reflect urine dilution; and (3) most OP pesticides have biological half-lives of only 12 to 36 hours (Needham 2005) and are metabolized and excreted from the body in hours to days (Barr 2008). Thus, OP pesticide metabolite excretion in a brief sampling period, whether measured in a spot, FMV, or 24-hour

urine sample, may not accurately represent a child's cumulative OP pesticide exposure over longer periods.

One recent study of 44 preschool-age children who provided between 10 and 26 biweekly spot urine samples over a 21-months reported that within-child variability for the five DAP metabolites measured exceeded between-child variability by several factors (Griffith et al. 2011), a finding which underscores concern about the reliability of this measure and raises questions about the source of within-child variability. To date, no studies have evaluated variability of OP metabolites within-day or in 24-hour urine samples collected several days apart. Nor have studies compared variability in DAP measurements when expressed in terms of unadjusted concentrations, creatinine-adjusted concentrations, or urinary excretion rates.

To address these questions, the objectives of this study with preschool-aged children in California were to: (1) evaluate the reproducibility of children's non-specific OP pesticide measurements within a one-week period; (2) evaluate the influence of concentration corrections (i.e., creatinine-adjustment and use of urinary excretion rate) on reproducibility of DAP measurements; (3) determine the degree of correspondence between pesticide metabolite concentration in spot urine samples (including FMVs) and same-day 24-hour samples; and (4) evaluate the sensitivity and specificity of spot urine samples to classify full-week average measurements.

Methods

Study population: We recruited a convenience sample of 25 children (10 boys, 15 girls) from clinics serving low-income families in the Salinas Valley, an agricultural region. Eligible children were between 3 and 6 years old, in good health with no history of diabetes or renal

disease, toilet-trained, free of enuresis, and had English- or Spanish-speaking mothers who were at least 18 years old. Sampling occurred in March and April 2004. The study was approved by the University of California at Berkeley Committee for the Protection of Human Subjects and parents provided written informed consent.

Urine sample collection: Each family participated for seven consecutive days (study days 1-7). On day 1, study staff provided families with sampling supplies and instructions on sample collection. Supplies included specimen trays (cleaned, sterile Specipan™, Baxter Scientific, McGaw Park, IL), gloves, collection jars with blank labels, a small refrigerator, and two 24-hour sampling record forms. On spot-sampling days (1, 3, 4, 6, and 7), families collected a single void at their convenience; families sometimes collected an FMV as the spot sample. On 24-hour sampling days (2 and 5), families were instructed to collect the child's FMV, all daytime and evening spot voids, and the FMV of the following day as separate specimens, and to note the timing of all voids (including missed voids) during this period on the 24-hr sampling record form.

Sample collection methods have been previously validated in this age range (Lu et al. 2001). Children voided directly into a collection jar or into a clean, sterile specimen tray placed on a training potty or toilet. If samples were collected in specimen trays, parents transferred specimens into collection jars. Parents identified each sample as an FMV or a non-FMV spot sample, recorded the time of collection on jar labels, and stored specimens in the refrigerator until daily collection by research staff. On 24-hour sampling days, research staff reviewed the sampling record form with the families to assure its accuracy and completeness.

Twenty-four-hour samples included three to eleven collected voids (mean 6.5). Overall, 86% (range=50-100%) of the reported voids during the 24 hour sampling periods were collected.

Twenty-two (44%) of the 24-hour samples were based on 100% collection of all voids. Reasons for missed voids included out-of-home bathroom use, toileting accidents, and participant error, such as missing an evening void. In sensitivity analyses, we excluded seven 24-hour samples with fewer than 5 total voids to confirm that collection errors did not substantially change study results (not shown).

Sample processing and analysis: Samples were processed at the Salinas Valley study office. The weight of each void was measured (grams) and the volume (milliliters) was estimated assuming a urine specific gravity of 1.022 (Salita et al. 1998). Individual voids from 24-hour sampling sessions which were not selected for individual analysis were pooled. After aliquotting, samples were stored at -80°C until shipped on dry ice to the Centers for Disease Control and Prevention for analysis.

We measured six DAP metabolites in all samples- three dimethyl (DM) phosphate metabolites: dimethylphosphate (DMP), dimethylthiophosphate (DMTP), and dimethyldithiophosphate (DMDTP); and three diethyl (DE) phosphates: diethylphosphate (DEP), diethylthiophosphate (DETP), and diethyldithiophosphate (DEDTP). These metabolites represent approximately 81% of the agricultural OP pesticides used in the Salinas Valley (Bradman et al. 2007). Laboratory methods and quality control procedures are described in detail elsewhere (Bravo et al. 2004). Briefly, samples were lyophilized to remove water, redissolved in a 1:1 solution of acetonitrile and diethyl ether, and analyzed using gas chromatography-tandem mass spectrometry using isotope dilution. Creatinine concentrations were determined using a commercially available method (Vitros CREA slides, Ortho Clinical Diagnostics, Raritan, NJ, USA). Westgard rules for quality control were used to establish the validity of each analytical

run (Westgard 2003). Limits of detection (LODs) were 0.2 µg/L for all DEs, 0.5 µg/L for DMP, 0.4 µg/L for DMTP, and 0.1 µg/L for DMDTP.

Levels below the LOD were imputed as $\text{LOD}/\sqrt{2}$ (Hornung and Reed 1990), and molar concentrations were summed within each sample to yield total DM, total DE, and total DAP concentrations. Metabolite levels in 24-hour voids were computed as the volume-weighted average of concentrations in all samples collected on sampling days 2 and 5, which included the FMV samples from the following day. We expressed metabolite concentration and/or excretion in three ways: (1) unadjusted metabolite concentration (nanomoles metabolite per liter of urine); (2) creatinine-adjusted metabolite concentration (nanomoles metabolite per gram of creatinine); and (3) urinary excretion rate (UER) of each metabolite (nanomoles metabolite per minute) computed as the total moles of excreted metabolite divided by the duration of time in minutes since the previous void. For 24-hour collections, time since the previous void was calculated as the time elapsed between the last void preceding the 24-hour sample collection (typically the last void on day 1 or 4) and the last void included in the 24-hour sample (typically the FMV on day 3 or 6).

Data Analysis

Statistical analyses were performed using Stata 10 for Windows (StataCorp LP, College Station, TX). We first computed descriptive statistics and assessed normality. Log_{10} transformed values were used for subsequent analyses.

Correspondence between spot samples and 24-hour samples: We constructed general estimating equation models using DAP concentration in a 24-hour sample as the outcome variable and concentration in a same-day spot sample (FMV or non-FMV) or a combination of these two spot samples as the predictor variable. Combinations of two spot samples were modeled as separate

predictor variables and as a single variable computed as either the arithmetic or volume-weighted average of the individual samples. Analyses using each DAP variable type (i.e., unadjusted concentrations, creatinine-adjusted concentrations, and UER) were compared. Robust standard errors were calculated.

Reproducibility of spot and 24-hour urine samples: We used mixed random effects models to compute the between-child and within-child variance for 24-hour samples from days 2 and 5, and for all FMV spot samples. For all non-FMV spot samples and for all spot samples (FMV and non-FMV), we estimated the variance attributable to each of three nested components – between-child, within-child, and within-child within-day variability (since multiple samples were available from the same day)– using two-level random-intercept models (Marchenko 2006). This method parses the total within-child variance into two parts; the component attributable to same-day within-child variability, and the component attributable to between-day within-child variability (Hauser et al. 2004; Preau et al. 2010; Ye et al. 2011). We calculated Akaike Information Criterion (AIC) values for the nested models with the largest number of observations (the all spot samples model) to compare the fit of models using unadjusted DAP concentrations, creatinine-adjusted concentrations, and UER. We also calculated the intraclass correlation coefficient (ICC) for each group of samples. The ICC is the ratio of between-subject variance to total variance.

Correlation of spot samples collected 0-6 days apart: We computed Pearson's correlations between pairs of samples collected on the same day (0 days apart) as well as on pairs of samples collected between 1 and 6 days apart. This analysis included all possible unique pairings of each child's spot samples, and was conducted separately for pairs of FMV spots only, pairs of non-

FMV spots only, and for all pairs of spots regardless of FMV status. Tests of significance were computed using a robust estimate of the variance.

Sensitivity and Specificity: We evaluated the sensitivity and specificity of one, two, or three randomly-selected spot urine samples from each child as predictors of high (top 20%) or elevated (top 40%) weekly average DAP metabolite concentrations. For “true” exposure, we calculated the arithmetic mean metabolite concentration of all spot urine samples collected from each child during the week, categorized these in quintiles, and assigned “true” high (top 20%) or elevated (top 40%) exposure level. For predictor sets, we created ten datasets each containing one randomly-selected spot sample per child (i.e. 25 observations total per dataset). Within each randomly-selected dataset, we again categorized the 25 metabolite concentration in quintiles, and assigned “predicted” high (top 20%) or elevated (top 40%) exposure levels. This process mimicked the exposure classification process in epidemiological studies, in which knowledge of the range and rankings of exposure in a population is often limited to what can be observed in the single sample or small number of samples collected per subject. The sensitivity and specificity figures we report represent the average sensitivity and specificity observed across the ten separate random samples. We report findings separately for FMV samples only, for non-FMV samples only, and for all spot samples. To determine whether collection of more than one spot sample could improve sensitivity, we repeated this analysis using the arithmetic mean of two or three spot samples from each child collected on different days. Because some children lacked three FMV or non-FMV samples from different days, we only assessed the sensitivity and specificity of three spot samples in the “Any Spot Sample” group.

Results

All children were Mexican-American and their ages ranged from 3 to 6.5 years (mean age: 4.5, SD: ± 0.93 years). Children urinated between 3 and 12 times (mean: 5.7 voids) per 24-hour period, and the volume of their individual spot samples ranged from 4.8 milliliters (mL) to 642.2 mL (mean: 146.6 mL) for FMV samples and from 8.4 to 238.3 mL (mean: 68.1 mL) for non-FMV spot samples. The creatinine concentration of spot samples ranged from 9.4 milligrams per deciliter (mg/dL) to 213.2 mg/dL (mean: 86.6 mg/dL) for FMV samples and from 4.5 to 158.7 mg/dL (mean: 68.6 mg/dL) for non-FMV samples. Creatinine concentration varied more within children than between children for both FMV (ICC=0.37) and non-FMV (ICC=0.22) spot samples.

Individual DE and DM metabolites were detected in >70% of samples, with the exception of DEDTP (17%) (data not shown). Total DM concentrations exceeded total DE concentrations, and total DAP levels predominantly reflected DM metabolites (Table 1). Central tendency measures (e.g. means, medians) for unadjusted concentration values were similar for FMV spot samples and 24-hour samples, which were generally higher than levels in non-FMV spot samples. In contrast, central tendency measures for creatinine-adjusted concentration values were more similar for FMVs and non-FMV spot samples, with both generally lower than creatinine-adjusted 24-hour samples.

Figure 1 presents the total DAP metabolite concentrations [in log10 scale (nanamoles per gram creatinine)] for each participant for all spot samples collected over the seven consecutive sampling days. There are no clear exposure trends over time among the children, though our data indicate that shifts of up to two orders of magnitude can occur over the week (see Figure 1, P7)

or within a single day (see Figure 1, P19, day 5). Twenty-four-hour samples collected only 3 days apart can likewise differ by an order of magnitude (e.g., Figure 1, P2, P7, P25).

Table 2 presents results of GEE models examining how well same-day spot samples predict 24-hour metabolite levels. For models examining a single unadjusted spot sample (FMV or non-FMV) and its respective 24-hour sample, the predictive power of the model, defined by the coefficient of determination (R^2), was highest for FMVs (R^2 for total DAP metabolites (nmol/L) = 0.54 for FMVs vs. 0.39 for non-FMV spots). The predictive power of the model was higher for creatinine-adjusted metabolites compared with unadjusted metabolites, especially for non-FMVs, with little difference between the sample types (total DAP R^2 =0.57 for FMV and 0.63 for non-FMVs). The predictive power was lowest for UER models.

The best fitting models were obtained when the arithmetic mean of an FMV and a non-FMV sample was used to predict the 24-hour values (Table 2). The best model fit was observed for creatinine-adjusted values (total DAP R^2 =0.84), but the variability in 24-hour metabolite levels explained by unadjusted values and UER estimates were comparable (R^2 =0.79 and 0.77, respectively). Model fit was strongest for analyses of total DM metabolites, and weakest for total DE metabolites. Volume-weighted averages of FMV and non-FMV spots produced model fits that were similar to that obtained with the arithmetic mean (data not shown). Results of sensitivity analyses which excluded seven 24-hour samples with fewer than 5 spot samples were consistent with those described above (data not shown).

AIC values for mixed random effects models used to estimate the variance in total DAPs attributable to between-child versus within-child sources based on all spot urine samples (N=247) indicated that best model fit was achieved with creatinine-adjusted values (AIC=364, Table 3) versus 372 and 451, respectively, for unadjusted concentration and UER models

(Supplemental Material, Table S1). Relative to unadjusted concentration and UER models, use of creatinine-adjusted values also provided the greatest degree of distinction between individual children [e.g. ICC= 0.32 for creatinine-adjusted total DAP metabolites in the “any spot sample” model (Table 3) versus ICC=0.21 and 0.15, respectively, for the analogous unadjusted concentration and UER models (Supplemental Material, Table S1)].

Among creatinine-adjusted total DAP concentration models (Table 3), non-FMV spot samples displayed the poorest reproducibility (ICC=0.27 for creatinine-adjusted total DAP metabolites). Inclusion of FMV samples in this analysis (the “Any Spot Sample” set) modestly increased the between-child relative to within-child variance (ICC=0.32) and suggested somewhat more between-day stability in measures (i.e. smaller within-child between-day variance). However, examination of 24-hour voids collected three days apart indicates that this between-day stability was limited, with 65% of total variance attributed to within-child variability (Table 3). A sensitivity analysis limited to the most complete 24-hour samples (based on 5 or more spot samples) indicated even greater within-child variance (for total DAPs, 77% of total, ICC=0.23; data not shown). The variance components of creatinine-adjusted total DM metabolites (Table 3) were similar to those for total DAP metabolites, though the reproducibility of DM samples was slightly higher (ICCs=0.30-0.39, compared with 0.27–0.35 for total DAPs). Reproducibility of creatinine-adjusted DE metabolites was much lower (ICCs=0.11-0.16) (Table 3), and within-child variance exceeded between-child variance between a factor between five (e.g., 57% versus 12% of the total variance for non-FMV spot samples) and eight (e.g., 88% versus 12% of the total variance for FMV spot samples) for these metabolites.

Based on their superior model performance in the preceding analyses, we used creatinine-adjusted values in subsequent analyses.

Table 4 presents Pearson correlations of creatinine-adjusted metabolite levels in spot samples collected 0-6 days apart. Correlations for total DAPs and total DMs were moderate (≈ 0.5) and statistically significant for samples collected on the same day (i.e. zero days apart) or one day apart, and became weaker as the number of days between samples increased. All correlations were weak and not significant for samples collected 5-6 days apart. DE metabolites had weaker correlations compared to DMs and the decay in correlations across days was more rapid for DEs compared to DM or total DAP metabolites. FMV samples tended to have higher correlations than non-FMV samples or any spot samples. The correlations between samples collected 2-3 days apart are comparable to Pearson correlations between levels in 24-hour samples collected 3 days apart, which were 0.35, 0.36, and 0.15, respectively, for total DAP, DM, and DE metabolites.

Table 5 presents results of sensitivity and specificity analyses. In this sample of participants, children with high (top 20%) weekly average total DAP metabolite concentrations would be correctly classified by a single non-FMV spot sample 52% of the time. Using two samples increased the sensitivity marginally to 56%. First morning voids also showed marginally higher sensitivity than non-FMV samples. Within the “any spot sample group”, three-spot predictors offered little apparent advantage over two-spot predictors. Overall, the sensitivity was higher to classify children in the top 40th percentile (elevated) than in the top 20th percentile, with a slight improvement with 2 versus 1 sample. Specificity was uniformly higher than sensitivity. Use of a single non-FMV spot sample would correctly identify a child in the lower 80% (i.e. not “high”) 88% of the time, and would identify a child in the lower 60th% (i.e. not “elevated”) 78% of the time.

Discussion

This study of variability in dialkylphosphate (DAP) metabolites of organophosphorous (OP) pesticides in urine samples from children 3-6 years old indicates high variability over a one week time-frame. Although we observed strong correlations between full-day 24-hour samples and same-day spot samples (see Pearson correlations, Table 2), there was weak correlation between DAP metabolite levels in 24-hour samples collected three days apart or in spot samples collected more than one day apart, with correlations weaker for diethyl (DE) compared to dimethyl (DM) DAP metabolites. Further, within-child variance was approximately two to three times the between-child variance for total DAP metabolites (e.g., 66% versus 34% for FMV samples for creatinine-adjusted total DAPs, ICC=0.27-0.35). This ratio was lower (and thus, the ICC higher) in dimethyl (DM) metabolites compared to diethyl (DE) metabolites (ICC=0.30-0.39 versus ICC=0.11-0.16, respectively). Finally, we found that DAPs measured in one, two, or even three spot urine samples have relatively low sensitivity to identify children who would be considered the most highly-exposed on the basis of their average full-week DAP concentrations. For example, a single non-FMV spot sample would correctly identify only 52% of the children whose true weekly exposure was in the top quintile, suggesting high Type II classification errors; by contrast, spot samples appear to offer good specificity, with a single non-FMV spot sample correctly identifying about 88% of children with true total DAP exposure below the top quintile. Overall, the high within-child variance, the weak correlation across days, and low specificity suggests that single day measurements may not adequately characterize exposure for longer time frames necessary for chronic risk assessments or epidemiologic studies.

Our analysis replicated methods applied by Ye (Ye et al. 2011), Preau (Preau et al. 2010), and colleagues to partition the variance components among between-child, within-child

between-day, and within-child within-day components for sample types with more than one sample per day (Table 3). Similar to their findings for bisphenol A and monoethyl phthalate, we found that overall within-person variance was higher than between-person variance, and that within-day variance was the largest component of total variance, suggesting that differences in exposure between days were lower in comparison with fluctuations that occurred during the course of each day. This finding suggests that eliminating within-day variability would substantially reduce within-child relative to between-child variability overall. However, we found that 65% of the total variance in 24-hour urine samples collected three days apart was due to within-child between-day variability, even though the 24 hour samples reflect complete sampling with no within-day variability (i.e., the entire sample was collected). These apparently contradictory findings suggest to us that the variance component analyses may overestimate the within-person within-day variance component. Specifically, we suspect the models may overestimate the degree of within-day variability that is biologically plausible, given that the total number of possible within-day observations is capped by the number of times individuals urinate each day, which in this study ranged between three and twelve times.

Our study provides important information about the appropriateness of creatinine adjustment to control for urinary dilution in children 3-6 years of age. Overall, our findings suggest that creatinine-adjustment of urinary DAP concentrations in children ages 3-6 years maximizes between-child relative to within-child variance (thus reducing exposure misclassification), improves estimation of the average 24-hour DAP concentration on the basis of spot urine samples, and may decrease the difference between FMV and easier-to-collect non-FMV spot samples in terms of their ability to estimate total 24-hour excretion. Despite this empirical evidence suggesting that creatinine adjustment effectively normalized the metabolite

concentrations, this method may introduce other sources of variability. Specifically, creatinine varies as a function of gender, age, meat consumption, body size, and muscle mass (Boeniger et al. 1993). Thus, large differences in size, muscle mass, and diet between developing children of the same age could result in very different normalized metabolite values among children with the same exposure. In future analyses, we will examine creatinine excretion patterns in these children and assess adjustment for specific gravity (not yet quantified in these samples) as an alternative approach to account for urinary dilution.

In summary, our findings raise concerns about the utility of urinary DAP metabolites as a biomarker of chronic OP pesticide exposure in young children. Two recent published analyses from our Center have detected no association between DAP measures in children and adverse outcomes, although they did detect significant associations between maternal prenatal DAP measurements and child neurodevelopmental outcomes (Bouchard et al. 2011; Marks et al. 2010), despite the fact that we have also observed high within versus between variability in maternal urine (Bradman et al. 2005). One interpretation of this finding is that OP exposures in childhood are less critical than exposures that occur *in utero*, i.e., the fetus is so exceptionally sensitive to OPs that the strength of the prenatal exposure effects are evident despite limitations in the exposure measure. The current study suggests another plausible explanation: that misclassification of child OP exposure biased results toward the null hypothesis.

Our study has several limitations. Collecting 24-hour urine samples from young children proved challenging, and missed voids due to occasional toileting accidents or other circumstances were beyond the control of study staff. Though 86% of the voids reported to have occurred during 24-hour sampling periods were collected, missed voids meant that individual spot samples made up a greater proportion of the 24-hour volume-weighted average than they

should have. Thus, our estimates of the association between metabolites in spot and 24-hour samples (Table 2) represent an upper bound. We measured class-specific organophosphate pesticide DAP metabolites, and our findings should not be generalized to pesticide-specific OP metabolites which may exhibit different variance patterns. Our study was also conducted in an agricultural community during the growing season, and it is possible that intermittent exposures related to nearby agricultural pesticide use may have caused higher variability in metabolite levels than would be observed in the general U.S. population. Furthermore, we note that different exposure scenarios could result in different variability patterns. Prior to the phase-out of diazinon and chlorpyrifos in household pesticides (U.S. EPA 2000, 2001), for example, children from homes with frequent indoor use of such pesticides may have shown more stable exposure levels. In a follow-up analysis, we intend to use additional data available to us to assess which potential exposure sources (e.g. diet, nearby agricultural pesticide use) best account for the metabolite variability patterns we present here.

Conclusions

We found that though DAP metabolites in single or multiple spot samples are strongly correlated with levels in same-day 24-hour samples, children's urinary DAP concentrations vary greatly from day to day, and use of spot samples to characterize a child's cumulative weekly exposure results in a moderate degree of misclassification. Exposure misclassification resulting from urinary metabolite variability has the potential to bias measures of association between early childhood OP exposures and developmental outcomes in epidemiologic research toward the null hypothesis; such misclassification may account for the weak or null findings often reported to date between pediatric urinary DAP concentrations and child development (Bouchard et al. 2011; Marks et al. 2010). Additional research on variability in measures of non-persistent

compounds in children is needed to assure that exposure biomarkers are valid and that epidemiologic studies have adequate power to detect health outcomes that may result from these exposures.

References

- Aprea C, Colosio C, Mammone T, Minoia C, Maroni M. 2002. Biological monitoring of pesticide exposure: a review of analytical methods. *J Chromatogr B Analyt Technol Biomed Life Sci* 769(2):191-219.
- Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL. 2005. Urinary creatinine concentrations in the U.S. population: implications for urinary biologic monitoring measurements. *Environ Health Perspect* 113(2):192-200.
- Barr DB. 2008. Biomonitoring of exposure to pesticides *Jourl of Chem Health and Safety* 15(6):20-29.
- Boeniger MF, Lowry LK, Rosenberg J. 1993. Interpretation of urine results used to assess chemical exposure with emphasis on creatinine adjustments: a review. *Am Ind Hyg Assoc J* 54(10):615-627.
- Bouchard MF, Chevrier J, Harley KG, Kogut K, Vedar M, Calderon N, et al. 2011. Prenatal Exposure to Organophosphate Pesticides and IQ in 7-Year-Old Children. *Environ Health Perspect* 119(8):1189-1195.
- Bradman A, Whitaker D, Quiros L, Castorina R, Claus Henn B, Nishioka M, et al. 2007. Pesticides and their metabolites in the homes and urine of farmworker children living in the Salinas Valley, CA. *J Expo Sci Environ Epidemiol* 17(4):331-349.
- Bradman A, Eskenazi B, Barr DB, Bravo R, Castorina R, Chevrier J, et al. 2005. Organophosphate urinary metabolite levels during pregnancy and after delivery in women living in an agricultural community. *Environ Health Perspect* 113(12):1802-1807.
- Bravo R, Calabiano LM, Weerasekera G, Whitehead RD, Fernandez C, Needham LL, et al. 2004. Measurement of dialkyl phosphate metabolites of organophosphorus pesticides in human

urine using lyophilization with gas chromatography-tandem mass spectrometry and isotope dilution quantification. *J Expo Anal Environ Epidemiol* 14(3):249-259.

Centers for Disease Control and Prevention. 2003. Second National Report on Human Exposure to Environmental Chemicals NCEH Pub. No. 02-0716. Atlanta, GA: Centers for Disease Control and Prevention, National Center Environmental Health.

Cornelis R, Heinzow B, Herber RF, Christensen JM, Poulsen OM, Sabbioni E, et al. 1996. Sample collection guidelines for trace elements in blood and urine. IUPAC Commission of Toxicology. *J Trace Elem Med Biol* 10(2):103-127.

Fenske RA, Lu C, Barr D, Needham L. 2002. Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. *Environ Health Perspect* 110(5):549-553.

Fenske RA, Bradman A, Whyatt RM, Wolff MS, Barr DB. 2005. Lessons learned for the assessment of children's pesticide exposure: critical sampling and analytical issues for future studies. *Environ Health Perspect* 113(10):1455-1462.

Fenske RA, Lu C, Simcox NJ, Loewenherz C, Touchstone J, Moate TF, et al. 2000. Strategies for assessing children's organophosphorus pesticide exposures in agricultural communities. *J Expo Anal Environ Epidemiol* 10(6 Pt 2):662-671.

Fenske RA, Elkner KP. 1990. Multi-route exposure assessment and biological monitoring of urban pesticide applicators during structural control treatments with chlorpyrifos. *Toxicol Ind Health* 6(3-4):349-371.

Griffith W, Curl CL, Fenske RA, Lu CA, Vigoren EM, Faustman EM. 2011. Organophosphate pesticide metabolite levels in pre-school children in an agricultural community: within- and between-child variability in a longitudinal study. *Environ Res* 111(6):751-756.

Hauser R, Meeker JD, Park S, Silva MJ, Calafat AM. 2004. Temporal variability of urinary phthalate metabolite levels in men of reproductive age. *Environ Health Perspect* 112(17):1734-1740.

Hornung RW, Reed LD. 1990. Estimation of Average Concentration in the Presence of Nondetectable Values. *Appl Occup Env Hyg* 5(1):46-51.

Kissel JC, Curl CL, Kedan G, Lu C, Griffith W, Barr DB, et al. 2005. Comparison of organophosphorus pesticide metabolite levels in single and multiple daily urine samples collected from preschool children in Washington State. *J Expo Anal Environ Epidemiol* 15(2):164-171.

Lu C, Knutson DE, Fisker-Andersen J, Fenske RA. 2001. Biological monitoring survey of organophosphorus pesticide exposure among pre-school children in the Seattle metropolitan area. *Environ Health Perspect* 109(3):299-303.

Marchenko Y. 2006. Estimating variance components in Stata. *The Stata Journal* 6(1):1-21.

Marks AR, Harley K, Bradman A, Kogut K, Barr DB, Johnson C, et al. 2010. Organophosphate pesticide exposure and attention in young Mexican-American children: the CHAMACOS study. *Environ Health Perspect* 118(12):1768-1774.

Martin MD, McCann T, Naleway C, Woods JS, Leroux BG, Bollen AM. 1996. The validity of spot urine samples for low-level occupational mercury exposure assessment and relationship to porphyrin and creatinine excretion rates. *J Pharmacol Exp Ther* 277(1):239-244.

Needham LL. 2005. Assessing exposure to organophosphorus pesticides by biomonitoring in epidemiologic studies of birth outcomes. *Environ Health Perspect* 113(4):494-498.

Preau JL, Jr., Wong LY, Silva MJ, Needham LL, Calafat AM. 2010. Variability over 1 week in the urinary concentrations of metabolites of diethyl phthalate and di(2-ethylhexyl) phthalate among eight adults: an observational study. *Environ Health Perspect* 118(12):1748-1754.

Salita M, Macknin M, Medendorp SV, Jahnke D. 1998. First-morning urine specific gravity and enuresis in preschool children. *Clin Pediatr (Phila)* 37(12):719-724.

U.S. EPA. 2000. Chlorpyrifos Revised Risk Assessment and Agreement with Registrants. Washington DC: U.S. Environmental Protection Agency.

U.S. EPA. 2001. Diazinon Revised Risk Assessment and Agreement with Registrants. Washington DC: U.S. Environmental Protection Agency.

Westgard JO. 2003. Westgard QC: Tools, Technology, and Training for Healthcare Laboratories. Available: <http://www.westgard.com> [accessed January 10 2003].

Ye X, Wong LY, Bishop AM, Calafat AM. 2011. Variability of urinary concentrations of bisphenol A in spot samples, first morning voids, and 24-hour collections. *Environ Health Perspect* 119(7):983-988.

Table 1. Unadjusted and creatinine-adjusted dialkylphosphate (DAP) concentrations in urine samples

<u>Type of Sample</u>	Unadjusted Concentration					Creatinine-Adjusted Concentration			
	<u>DF^a(%)</u>	<u>GM^b</u>	<u>Mean</u>	<u>Median</u>	<u>Max</u>	<u>GM^b</u>	<u>Mean</u>	<u>Median</u>	<u>Max</u>
Non-FMV Spot Samples (n=137)									
Total DAPs	99.3	110	239	122	4820	196	529	218	20600
Total DMs	94.2	63.0	179	60.3	4790	112	436	96.2	20500
Total DEs	95.6	26.8	59.8	31.1	500	47.7	98.8	68.0	529
First Morning Void (FMV) Samples (n=110)									
Total DAPs	98.2	162	307	157	2530	205	483	200	11800
Total DMs	95.5	92.3	234	94.4	2380	117	392	124	11600
Total DEs	97.3	43.4	72.5	57.1	391	55.1	91.1	77.6	376
24-Hour Samples (n=50)									
Total DAPs	--	158	296	144	3700	275	621	245	10100
Total DMs	--	94.6	230	89.9	3590	166	508	139	9920
Total DEs	--	45.9	65.0	53.3	248	78.9	113	93.6	610

^aDetection frequency. ^bGeometric mean.

Table 2. Modelling of 24-hour metabolite excretion^a using same-day spot urine samples as predictors (N=41^b)

Type of Spot Sample	Metabolite Excretion Units ^c	Metabolite Type	Regression Model				Pearson Correlation	
			β	(95% CI)	Intercept	Model R ²	r	P-value
Non-FMV Spot	Unadjusted (nmol/L)	Total DAPs	0.52	(0.28,0.76)	1.11	0.39	0.63	<0.001
		Total DMs	0.60	(0.41,0.78)	0.86	0.49	0.70	<0.001
		Total DEs	0.25	(0.06,0.43)	1.30	0.14	0.37	0.011
	Creatinine-Adjusted (nmol/g creat)	Total DAPs	0.66	(0.48,0.84)	0.87	0.63	0.80	<0.001
		Total DMs	0.69	(0.53,0.84)	0.74	0.68	0.82	<0.001
		Total DEs	0.33	(0.16,0.50)	1.31	0.25	0.50	<0.001
	Urinary Excretion Rate (nmol/min)	Total DAPs	0.51	(0.23,0.79)	-0.69	0.42	0.65	0.001
		Total DMs	0.57	(0.37,0.77)	-0.72	0.52	0.72	<0.001
		Total DEs	0.21	(0.01,0.40)	-1.49	0.09	0.30	0.036
First Morning Void (FMV) Spot	Unadjusted (nmol/L)	Total DAPs	0.66	(0.42,0.90)	0.70	0.54	0.74	<0.001
		Total DMs	0.66	(0.42,0.89)	0.66	0.57	0.75	<0.001
		Total DEs	0.39	(0.18,0.60)	0.99	0.26	0.51	0.001
	Creatinine-Adjusted (nmol/g creat)	Total DAPs	0.75	(0.50,1.01)	0.68	0.57	0.76	<0.001
		Total DMs	0.72	(0.47,0.97)	0.71	0.59	0.77	<0.001
		Total DEs	0.41	(0.19,0.63)	1.15	0.27	0.52	0.001
	Urinary Excretion Rate (nmol/min)	Total DAPs	0.43	(0.13,0.74)	-0.72	0.37	0.60	0.008
		Total DMs	0.51	(0.22,0.79)	-0.71	0.45	0.67	0.001
		Total DEs	0.26	(0.04,0.49)	-1.36	0.19	0.43	0.025
Average of Non-FMV and FMV Spots ^d	Unadjusted (nmol/L)	Total DAPs	0.90	(0.76,1.05)	0.17	0.79	0.89	<0.001
		Total DMs	0.92	(0.78,1.07)	0.09	0.84	0.92	<0.001
		Total DEs	0.66	(0.78,1.07)	0.57	0.44	0.66	<0.001
	Creatinine-Adjusted (nmol/g creat)	Total DAPs	0.92	(0.81,1.03)	0.20	0.84	0.92	<0.001
		Total DMs	0.93	(0.81,1.04)	0.14	0.89	0.94	<0.001
		Total DEs	0.70	(0.44,0.96)	0.60	0.48	0.69	<0.001
	Urinary Excretion Rate (nmol/min)	Total DAPs	0.83	(0.64,1.03)	-0.30	0.77	0.88	<0.001
		Total DMs	0.85	(0.69,1.01)	-0.32	0.82	0.91	<0.001
		Total DEs	0.66	(0.34,0.99)	-0.67	0.39	0.62	<0.001

Table 2 (Continued)

^aSpot and 24-hr samples analyzed on \log_{10} scale. ^bN=41 child-days; 9 samples lacked an FMV sample and were excluded. Regression models are clustered by child. ^cUnits of both the spot sample(s) and the 24-hour sample. ^dValues converted to \log_{10} scale after arithmetic average of concentrations on normal scale was computed; not volume-weighted.

Table 3. Variance apportionment of log-transformed creatinine-adjusted DAP metabolite concentrations in spot urine samples collected during one week and in 24-hour voids collected 3 days apart (N=25)

Type of Sample	N ^a	Total DAPs			Total DMs			Total DEs		
		Variance	% Total Variance	ICC	Variance	% Total Variance	ICC	Variance	% Total Variance	ICC
Non-FMV Spot Samples										
Between child	137	0.075	25%	0.27 ^b	0.114	27%	0.30 ^b	0.041	12%	0.14 ^b
Within child, between day		0.105	34%		0.149	35%		0.108	31%	
Within child, within day		0.125	41%		0.162	38%		0.198	57%	
First Morning Void (FMV) Samples										
Between child	110	0.105	34%	0.34	0.172	38%	0.39	0.035	12%	0.11
Within child ^c		0.207	66%		0.276	62%		0.246	88%	
Any Spot Samples (FMV or Non-FMV)										
Between child	247	0.097	31%	0.32 ^b	0.157	36%	0.37 ^b	0.033	10%	0.11 ^b
Within child, between day		0.064	20%		0.070	16%		0.049	15%	
Within child, within day		0.152	49%		0.213	48%		0.238	74%	
24-Hour Voids										
Between child	50	0.089	35%	0.35	0.131	36%		0.026	16%	0.16
Within child ^c		0.168	65%		0.237	64%		0.132	84%	

^a Number of samples used in calculation. ^bThe ICC presented is the ratio of between-child to total variability as calculated using a one-factor (child) as opposed to a two-factor nested mixed-effects model. ^cBecause FMV spots and 24-hour voids allow only one measure per day, the distinction between within-day versus between-day variability is not applicable.

Table 4. Pearson correlations of creatinine-adjusted DAP metabolite concentrations (log₁₀ scale) in paired, same-child spot urine samples collected zero to six days apart

Type of Spot Sample MetaboliteType	Days Elapsed Between Paired Samples						
	0 (Same Day)	1	2	3	4	5	6
Non-FMV Spot Samples	(N=26)	(84)	(60)	(58)	(65)	(24)	(2)^b
Total DAPs	0.54*	0.48*	0.08	0.24	0.16	-0.01	—
Total DMs	0.58*	0.45*	0.27*	0.37*	0.10	-0.15	—
Total DEs	0.44*	0.34*	-0.21	0.13	0.13	0.31	—
First MorningVoid (FMV) Samples	(N=0)	(58)	(57)	(45)	(20)	(21)	(2)^b
Total DAPs	—	0.54*	0.27*	0.27	0.26	0.25	—
Total DMs	—	0.59*	0.34*	0.30*	0.38	0.31	—
Total DEs	—	0.25	-0.01	0.32*	-0.10	-0.05	—
Any Spot Samples (FMV or Non-FMV)	(N=92^a)	(303)	(248)	(203)	(154)	(77)	(25)
Total DAPs	0.46*	0.45*	0.25*	0.23*	0.17*	0.16	0.16
Total DMs	0.49*	0.48*	0.35*	0.30*	0.21*	0.18	0.13
Total DEs	0.25*	0.22*	-0.15	0.08	0.02	0.11	-0.27

Note: * signifies p-value ≤ 0.05 . ^aN refers to unique, within-child pairings of spot samples which meet pairing requirements (e.g. N=92 discrete pairings of within-child spots collected on the same day, or 0 days apart). ^bNot reported due to small sample size.

Table 5. Sensitivity and specificity^a for classifying children with high (top 20%) and elevated (top 40%) one-week average total DAP metabolite concentrations^b with one, two, or three urine samples

Type of Sample	High (Top 20%)		Elevated (Top 40%)	
	Sensitivity	Specificity	Sensitivity	Specificity
Non-FMV Spot Samples				
One Sample	0.52	0.88	0.67	0.78
Two Samples ^c	0.56	0.89	0.68	0.79
First MorningVoid (FMV) Samples				
One Sample	0.58	0.90	0.65	0.77
Two Samples ^c	0.58	0.90	0.78	0.85
Any Spot Samples (FMV or Non-FMV)				
One Sample	0.46	0.87	0.63	0.75
Two Samples ^c	0.60	0.90	0.76	0.84
Three Samples ^c	0.64	0.91	0.73	0.82

^a Figures presented reflect the average sensitivity and specificity calculated based on ten separate random samples of predictor spot samples. ^b Calculations use creatinine-adjusted total DAP metabolite concentration (nmol/gram creatinine) on the log₁₀ scale. ^c Pairs and trios consist of spot samples collected on different days (i.e., no pairs of same-day spots).

Figure Legend

Figure 1. Concentration of total DAP metabolites (nmol/ gram creatinine) in log₁₀ scale for all spot samples and 24-hour samples collected over one week. Each panel represents an individual participant (labeled as P1 – P25.) The dots in each panel represent the total DAP metabolite concentration in the 24-hour samples from sampling days 2 and 5, respectively. The panels are ranked in order of ascending arithmetic mean of all spot samples collected for each child. Thus, the bottom row (P21-P25) contains the children rated as having “true” high (top 20%) weekly exposure in the sensitivity and specificity analysis, and the bottom two rows (P16-P25) contain the children with “true” elevated (top 40%) weekly exposure.

